

A Legendrian surgery presentation of contact 3-manifolds

Fan Ding and Hansjörg Geiges

Abstract

We prove that every closed, connected contact 3-manifold can be obtained from S^3 with its standard contact structure by contact (± 1) -surgery along a Legendrian link. As a corollary, we derive a result of Etnyre and Honda about symplectic cobordisms (in slightly stronger form).

Mathematics Subject Classification (2000). Primary 53D35; Secondary 57R65, 57R90.

Keywords. Contact structure, Legendrian link, contact surgery, symplectic cobordism.

1 Introduction

A *contact structure* ξ on a differential 3-manifold M is a smooth tangent 2-plane field such that any differential 1-form α that locally defines ξ as $\xi = \ker \alpha$ satisfies the nonintegrability condition $\alpha \wedge d\alpha \neq 0$. Notice that the sign of $\alpha \wedge d\alpha$ only depends on ξ , not the choice of α . In particular, M has to be orientable. Without any essential loss of generality, we assume our 3-manifolds to be *oriented*, and we restrict attention to *positive* contact structures, defined by the requirement that $\alpha \wedge d\alpha$ be a *positive* volume form. Moreover, our contact structures will be understood to be *coorientable*, which is equivalent to saying that a defining 1-form α exists globally.

The *standard contact structure* ξ_0 on $S^3 \subset \mathbb{R}^4$ is defined by $\sum_{i=1}^2 (x_i dy_i - y_i dx_i) = 0$. A theorem of Lutz [19] and Martinet [21] states that on any given (closed, orientable) 3-manifold M one can find a contact structure in each homotopy class of 2-plane fields by performing surgery on (S^3, ξ_0) along a link *transverse* to ξ_0 . (The part about ‘each homotopy class of 2-plane fields’ is not completely covered by the cited references, but belongs to contact geometric folklore, cf. [9].)

Recall the dichotomy between *overtwisted* and *tight* contact structures. A contact structure ξ on M is called overtwisted if one can find an embedded disc $D \hookrightarrow M$ with boundary ∂D tangent to ξ , but D transverse to ξ along ∂D . If no such D exists, then ξ is called tight. Eliashberg has shown that the (isotopy) classification of overtwisted contact structures coincides with the (homotopy) classification of tangent 2-plane fields [3].

In the Lutz-Martinet theorem no statement is made whether *every* contact structure on M can be obtained via transverse contact surgery as described, but from the work of Eliashberg [3] one can deduce that at least all overtwisted contact structures (up to isotopy) are covered by this construction.

A smooth knot K in a contact manifold (M, ξ) is called *Legendrian* if it is everywhere tangent to ξ . Such a Legendrian knot inherits a canonical *contact framing* of its normal bundle, defined via any vector field in $\xi|_K$ transverse to K . In [2] we described a notion of contact r -surgery along such Legendrian knots, where the surgery coefficient $r \in \mathbf{Q} \cup \{\infty\}$ is measured with respect to the contact framing; details will be recalled below. In general the resulting contact structure on the surgered manifold depends on choices, but for $r = 1/k$ with $k \in \mathbb{Z}$ it is fully determined by (M, ξ) , K and r [2, Prop. 7] (up to contactomorphism, i.e. diffeomorphism preserving the contact structures). The inverse of a contact $(1/k)$ -surgery is a contact $(-1/k)$ -surgery; for the case $k = \pm 1$ (which is the one most relevant to our discussion) we give an explicit proof of this fact in Section 3.

Our main theorem is the following.

Theorem 1. *Let (M, ξ) be a closed, connected contact 3-manifold. Then (M, ξ) can be obtained by contact (± 1) -surgery along a Legendrian link in (S^3, ξ_0) .*

Contact (-1) -surgery coincides with the symplectic handlebody surgery described by Eliashberg [4] and Weinstein [25], cf. [15] and Section 3 below. In particular, if (M', ξ') is obtained from (M, ξ) by contact (-1) -surgery along a Legendrian link, then there is a so-called *Stein cobordism* from (M, ξ) to (M', ξ') , see [7] for details, and [6] for background information. This is denoted by $(M, \xi) \prec (M', \xi')$. If $(S^3, \xi_0) \prec (M', \xi')$, then (M', ξ') is *Stein fillable*; the concave end (S^3, ξ_0) of the Stein cobordism can be filled with the standard symplectic 4-ball.

It would be possible to deduce Theorem 1 directly from the results of Eliashberg [4]. By two contact $(+1)$ -surgeries on a given (M, ξ) one can effect a simple Lutz twist (Proposition 9 of the present paper). This produces an overtwisted contact structure on M . Surgery along a suitable link in M produces S^3 . Eliashberg's results, in particular [4, Lemma 2.4.3], then say that S^3 (with some contact structure determined by the choice of surgeries) can be obtained from M by contact (-1) -surgeries. (Alternatively, one may appeal to [15, Lemma 4.4] and then directly to [4, Thm. 1.3.4].) As shown in Section 6 below, a further sequence of contact (-1) -surgeries on this S^3 will produce (S^3, ξ_0) .

A. Stipsicz [23] has suggested a similar shortcut to the proof of our theorem.

The extra benefit of the proof of Theorem 1 given in the present paper lies in the fact that it contains an algorithm for turning contact surgeries with rational coefficients into surgeries of the desired kind. It thus allows to discuss explicit surgery descriptions of contact manifolds. We hope to elaborate on this point in a future publication.

It is now an easy matter to derive from Theorem 1 the following corollary, which generalises a theorem of Etnyre and Honda [7, Thm. 1.1].

Corollary 2. *Let (M_i, ξ_i) be contact 3-manifolds (as in Theorem 1), $i = 1, \dots, n$. Then there is a Stein fillable contact manifold (M, ξ) such that $(M_i, \xi_i) \prec (M, \xi)$ for each i .*

Proof. Let (M_i, ξ_i) be obtained from (S^3, ξ_0) by contact $(+1)$ -surgery along a link L_i^+ and (-1) -surgery along a link L_i^- . We may assume that the links $L_1^+, L_1^-, \dots, L_n^+, L_n^-$ are pairwise disjoint. Let (M, ξ) be the contact manifold obtained from (S^3, ξ_0) by contact (-1) -surgery along the link $L_1^- \cup \dots \cup L_n^-$. Then (M, ξ) is Stein fillable. Moreover, each (M_i, ξ_i) is obtained from (M, ξ) by contact $(+1)$ -surgeries, so (M, ξ) is obtained from (M_i, ξ_i) by contact (-1) -surgeries.

Etnyre and Honda, by contrast, base their proof on a result of Giroux [13], [14] about open book decompositions adapted to contact structures. In principle, their methods suffice to deduce this stronger corollary, too. Our proof arguably has the advantage that a surgery description is more appropriate in the context of a cobordism theoretic result than an open book decomposition.

2 Contact surgery

We recall a few definitions and results from [2], chiefly to fix notation and conventions.

Let K be a Legendrian knot in a contact 3-manifold (M, ξ) . *Rational surgery* on K with coefficient $r = p/q \in \mathbf{Q} \cup \{\infty\}$ (with p, q coprime) is defined as follows: Denote a tubular neighbourhood of K (diffeomorphic to a solid torus) by νK . Let (μ, λ) be a positively oriented basis for $H_1(\partial\nu K; \mathbf{Z}) \cong \mathbf{Z} \oplus \mathbf{Z}$, where λ is determined up to sign as the class of a parallel copy of K determined by the contact framing, and μ is determined by a suitably oriented meridian (i.e. a nullhomologous circle in νK), cf. [15, p. 672]. We obtain a new manifold M' by cutting νK out of M and regluing it by a diffeomorphism of $\partial(\nu K)$ sending μ (on the boundary of the solid torus to be glued in) to the curve $p\mu + q\lambda$ (on the boundary of $M - \nu K$). This procedure determines M' up to orientation-preserving diffeomorphism. For $r = \infty$ the surgery is trivial.

There is a unique contact geometric model for the tubular neighbourhood of a Legendrian knot. In order to describe this, we consider $N = \mathbf{R}^2 \times (\mathbf{R}/\mathbf{Z})$ with coordinates (x, y, z) and contact structure ζ defined by

$$\cos(2\pi z) dx - \sin(2\pi z) dy = 0.$$

For each $\delta > 0$, let

$$N_\delta = \{(x, y, z) \in N : x^2 + y^2 \leq \delta^2\}.$$

We identify $T_\delta = \partial N_\delta$ with $\mathbf{R}^2/\mathbf{Z}^2$, using the contact framing, and write (μ, λ) for a positively oriented basis for $H_1(T_\delta; \mathbf{Z}) \cong \mathbf{Z} \oplus \mathbf{Z}$, with μ corresponding to a meridian and λ to a longitude determined by this framing. A possible representative of λ would be

$$\{(\delta \sin(2\pi z), \delta \cos(2\pi z), z) : z \in \mathbf{R}/\mathbf{Z}\}.$$

The vector field $X = x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y}$ is a *contact vector field* for the contact structure ζ , i.e. its flow preserves ζ . This vector field is transverse to T_δ . Any closed surface in a contact manifold that admits a transverse contact vector field is called *convex* [10].

The set Γ_{T_δ} of points $w \in T_\delta$ where $X(w) \in \zeta(w)$ is equal to

$$\Gamma_{T_\delta} = \{(\pm\delta \sin(2\pi z), \pm\delta \cos(2\pi z), z) : z \in \mathbf{R}/\mathbf{Z}\}.$$

This set is called the *dividing set* of T_δ ; its isotopy type is independent of the choice of contact vector field transverse to T_δ . Notice that the dividing set is made up of two copies of the longitude determined by the contact framing.

In general, if T is a convex torus in a tight contact 3-manifold, the dividing set Γ_T consists of an even number $\#\Gamma_T$ of parallel essential curves. (Conversely, the absence of homotopically trivial curves on a convex surface $\Sigma \neq S^2$ in a contact manifold guarantees that Σ has a tight neighbourhood.) This is (a special case of) Giroux's criterion, cf. [16, Thm. 3.5]. After a diffeomorphism isotopic to the identity, one may assume the dividing curves to be linear relative to an identification of T with $\mathbf{R}^2/\mathbf{Z}^2$. Their slope will be called the *slope* $s(T)$ of the convex torus T . If T bounds a solid torus, our convention will be to identify it with $\mathbf{R}^2/\mathbf{Z}^2$ in such a way that the meridian μ has slope 0 and the (chosen) longitude λ has slope ∞ . Thus, with respect to the contact framing, the T_δ above is a convex torus with $\#\Gamma_{T_\delta} = 2$ and $s(T_\delta) = \infty$. The key observation by Giroux is that the dividing set of a convex surface Σ encodes all the essential contact geometric information in a neighbourhood of Σ . In particular, one can glue contact manifolds along surfaces with the same dividing set (taking boundary orientations into account). The specific results of Giroux necessary to make this statement precise are summarised in Section 2 of [2].

We now describe how to perform this rational surgery on K in such a way that the resulting manifold again carries a contact structure. Write

$$C = \{(x, y, z) \in N : x = y = 0\}$$

for the spine of N . Then there is a contact embedding $f : (N_2, \zeta) \rightarrow (M, \xi)$ such that $f(C) = K$. We want to construct a contact structure ξ' on the manifold M' obtained from M by rational surgery on K with coefficient $r = p/q \in \mathbf{Q} \cup \{\infty\}$. Let

$$P = \{(x, y, z) \in N : 1 \leq x^2 + y^2 \leq 4\} = N_2 \setminus \text{Int } N_1.$$

Let $g : P \rightarrow P$ be an orientation-preserving diffeomorphism sending T_δ to T_δ , $\delta = 1, 2$, and μ to $p\mu + q\lambda$.

If $p \neq 0$, then $(g_*)^{-1}(\zeta)$ is a contact structure on P with respect to which T_δ is a convex torus of non-zero slope. By [16, Thm. 2.3], which gives an enumeration of tight contact structures on the solid torus with convex boundary as in our situation (and in particular shows this set of contact structures to be non-empty), the contact structure $(g_*)^{-1}(\zeta)$ on P can be extended to a tight contact structure ζ' on N_2 . Define

$$M' = (M - f(N_1)) \cup N_2 / \sim,$$

where $w \in P \subset N_2$ is identified with $f(g(w)) \in M$, and equip M' with the contact structure ξ' it inherits from (M, ξ) and (N_2, ζ') . We say that (M', ξ') is obtained from (M, ξ) by *contact r -surgery on K , $r \neq 0$* .

Topologically, M' is completely determined by K and r , but ξ' depends on the choice of ζ' . Only for $r = 1/k$, $k \in \mathbf{Z}$, this choice is unique. Moreover, as pointed out in the introduction, contact $(-1/k)$ -surgery is the inverse of contact $(1/k)$ -surgery [2, Prop. 8]; see the following section for the case $k = \pm 1$.

If $p = 0$ (and $q = \pm 1$), then $(g_*)^{-1}(\zeta)$ will be a contact structure on P with respect to which T_δ is a convex torus of slope zero. In that case we use a construction that will appear again in various guises in the discussion of Lutz twists below. Namely, consider a solid torus $S^1 \times D^2$ with S^1 -coordinate θ and polar coordinates r, φ on D^2 . Write μ_0 for the meridian of this solid torus and λ_0 for a longitude given by setting φ equal to a constant. Let ζ_0 be the contact structure on $S^1 \times D^2$ defined by the differential 1-form $\beta_0 = h_1(r) d\theta + h_2(r) d\varphi$. Here we impose the following general conditions on the smooth plane curve $r \mapsto \gamma(r) = (h_1(r), h_2(r))$:

- $h_1(r) \equiv \pm 1$ and $h_2(r) = \pm r^2$ for r near 0; this guarantees that the 1-form β_0 is actually defined at $r = 0$.
- Position vector $\gamma(r)$ and velocity vector $\gamma'(r)$ are never parallel to each other (in particular $\gamma(r) \neq (0, 0)$); this ensures that $\beta_0 = 0$ does indeed define a contact structure, cf. [8].

Without loss of generality we shall always assume that $\gamma(r)$ winds in counter-clockwise (i.e. positive) direction around the origin. This amounts to fixing the orientation of $S^1 \times D^2$ as the one given by $d\theta \wedge r dr \wedge d\varphi$.

In order to define 0-surgery, we further require that $h_1(1) = 1$ and $h_2(1) = 0$, and that $\gamma(r)$ completes exactly one half-turn as r increases from 0 to 1. Then along the boundary $T^2 = \partial(S^1 \times D^2)$ the contact structure is given by $d\theta = 0$. Recall that the *characteristic foliation* Σ_ξ of a surface Σ in a contact manifold (M, ξ) is the singular 1-dimensional foliation defined by the intersection of the tangent planes of Σ with ξ . Hence, the characteristic foliation $T_{\zeta_0}^2$ consists of the meridional loops. By a C^∞ -small perturbation of T^2 (inside a slight thickening of $S^1 \times D^2$) one can turn this 2-torus into a convex surface with $\# \Gamma_{T^2} = 2$ and slope 0, see [11, p. 795/6]. Then the solid torus with this convex boundary can be glued into $M - f(N_1)$, using the attaching map corresponding to $p = 0$, i.e. the map that sends μ_0 to λ and λ_0 to $-\mu$, say. (Notice, in particular, that the proviso ‘if such exists’ in Remark (1), Section 3 of [2] was superfluous.)

3 Contact (± 1) -surgeries

We follow Weinstein [25] in the description of contact (± 1) -surgeries as symplectic handlebody surgeries. Consider $\mathbb{R}^4$ with coordinates x, y, z, t and standard symplectic form

$\omega = dx \wedge dy + dz \wedge dt$. The vector field

$$Z = 2x\partial_x - y\partial_y + 2z\partial_z - t\partial_t$$

is a Liouville vector field for ω , that is,

$$L_Z\omega = d(i_Z\omega) = \omega.$$

This implies that $\alpha = i_Z\omega$ defines a contact form on any hypersurface in $\mathbb{R}^4$ transverse to Z .

First we consider the hypersurface $V = \{y^2 + t^2 = \varepsilon^2\}$. Introduce a new coordinate θ by setting $y = \varepsilon \cos \theta$, $t = \varepsilon \sin \theta$. Then

$$\alpha|_V = \varepsilon((-2x \sin \theta + 2z \cos \theta) d\theta + \cos \theta dx + \sin \theta dz).$$

Observe that the circle $L = V \cap \{x = z = 0\}$ is Legendrian, and that the radial vector field $x\partial_x + z\partial_z$ is a contact vector field. The contact structure near L is not quite in the normal form considered in the previous section, but it is sufficient to know that small enough neighbourhoods of any two Legendrian knots are contactomorphic.




Figure 1: Contact (-1) -surgery.

Figure 1 shows how to attach a 2-handle (with boundary transverse to Z) along a tubular neighbourhood of L in V . We want to show that this amounts to a contact (-1) -surgery along L : The orientation on V is given by $\alpha \wedge d\alpha = \varepsilon^2 dx \wedge dz \wedge d\theta$, and the contact framing of L in V by $\sin \theta \partial_x - \cos \theta \partial_z$. This implies that the framing of the surgery, given by ∂_x , makes one negative twist in the xz -plane relative to the contact framing as we go once around L in θ -direction (this sign is independent of the choice of orientation for L).

Given a Legendrian knot K in a contact manifold (M, ξ) , a small neighbourhood of K is contactomorphic to a neighbourhood of L in V . Therefore the handle described in Figure 1 can be used to perform a contact (-1) -surgery on K .

If (M, ξ) is strongly symplectically fillable ($\emptyset \prec (M, \xi)$ in the notation of the introduction), that is, M is the boundary of a symplectic manifold (W, ω) admitting a Liouville vector field Z near $\partial W = M$, transverse to M , pointing outwards, and satisfying $\ker i_Z \omega = \xi$, then the construction shows that the surgered manifold will again be strongly symplectically fillable (for this one also needs to ‘glue’ the Liouville vector field on W near $K \subset M$ and that in the handle picture near L , see [25] for the technical details).

On $V' = \{x^2 + z^2 = \delta^2\}$ by contrast, and with $x = \delta \cos \varphi$, $z = \delta \sin \varphi$, we have

$$\alpha|_{V'} = \delta((-y \sin \varphi d\varphi + t \cos \varphi d\varphi) + 2 \cos \varphi dy + 2 \sin \varphi dt).$$

Now we have $\alpha \wedge d\alpha = 2\delta^2 dt \wedge dy \wedge d\varphi$, and the contact framing of $L' = V' \cap \{y = t = 0\}$ is given by $\sin \varphi \partial_y - \cos \varphi \partial_t$. So the framing ∂_y of the surgery indicated in Figure 2 makes one positive twist in the ty -plane relative to the contact framing when going once around L' in φ -direction. Hence this amounts to a contact $(+1)$ -surgery. It can be performed as a symplectic handlebody surgery on the concave end of a symplectic cobordism (where the Liouville vector field points inward).




Figure 2: Contact $(+1)$ -surgery.

Figure 3 shows that the composition of a contact (-1) -surgery and a contact $(+1)$ -surgery leads to a manifold contactomorphic to the one we started from. Given two hypersurfaces in a symplectic manifold, both transverse to the same Liouville vector field Z and with contact structures induced by Z , one can show by a straightforward calculation (cf. [25, Lemma 2.2]) that the local diffeomorphism between the two hypersurfaces

defined by following the integral curves of Z is a contactomorphism. Thus, we simply have to move the ‘bump’ in Figure 3 along the integral curves of Z back to the original hypersurface V .




Figure 3: Composition of a contact (-1) -surgery and a $(+1)$ -surgery.

The same figure gives an elementary proof (not appealing to an h -principle, for instance) that an iteration of contact (± 1) -surgeries may be replaced by a simultaneous contact surgery (without changing the surgery coefficients) along a suitable link. This amounts, as it were, to a handle reordering lemma for contact surgeries. Suppose we have performed a surgery along a Legendrian knot K_1 as in Figure 1, and we now want to carry out a further surgery along a Legendrian knot K_2 in the surgered manifold. By a contact isotopy we can move K_2 away from the belt sphere of the first surgery (such contact isotopies can easily be constructed using contact Hamiltonians, cf. [9]). As before, we see that the complement of a neighbourhood of this belt sphere is contactomorphic to the complement of a neighbourhood of K_1 in the original manifold. Using the radial contact vector field near K_1 , we can move K_2 away from the first handle.

4 Front projections

As a first preparation for the proof of Theorem 1 we recall how to describe Legendrian links in $\mathbf{R}^3$ with its standard tight contact structure $dz + x dy = 0$ (which is contactomorphic to (S^3, ξ_0) with a point removed, and the universal local model for contact structures).

We visualise links in $\mathbf{R}^3$ by projecting them into the (y, z) -plane. The condition for $p(t) = (x(t), y(t), z(t))$ to describe a Legendrian curve in $\mathbf{R}^3$ is that $z'(t) + x(t)y'(t) = 0$. Any closed curve $\bar{p}(t) = (y(t), z(t))$ in the (y, z) -plane that may have cusps and transverse self-crossings, but does not have any vertical tangencies (such a curve will be called a *front*), comes from a unique Legendrian knot $p(t)$ in $\mathbf{R}^3$, found by setting $-x(t)$ equal to the slope of $\bar{p}(t)$. Cusps of $\bar{p}(t)$ correspond to points where $p(t)$ is parallel to the x -axis.

The plane curve $\bar{p}(t)$ is called the *front projection* of $p(t)$. The theory of front projections was developed by Arnol'd [1]; all information relevant for our purposes can be found in Section 1 of [15].

There are two Legendrian isotopy invariants of oriented Legendrian knots K , viz. the Thurston-Bennequin invariant $tb(K)$ and the rotation number $r(K)$. The Thurston-Bennequin invariant measures the linking number of K with its push-off determined by the contact framing; we refer to [15] for the definition of the rotation number. In the present context we only need to know the following properties of these invariants:

- If the front projection $\bar{K}$ of the Legendrian knot $K \subset \mathbf{R}^3$ has no crossings, then $tb(K) = -c(\bar{K})/2$, where $c(\bar{K})$ denotes the number of cusps of the front projection. (This is immediate from the definition of $tb(K)$.)
- The rotation number $r(K)$ is equal to $\lambda_-(\bar{K}) - \rho_+(\bar{K})$, where λ_- denotes the number of cusps with vertex on the left and oriented downwards, ρ_+ the number of right cusps oriented upwards (remember that K has to be oriented). Changing the orientation of K will change the sign of $r(K)$ (and leave $tb(K)$ unaltered).
- For the trivial knot K , these invariants satisfy the Thurston-Bennequin inequality

$$tb(K) + |r(K)| \leq -1.$$

The consequences of these properties that we shall use in the next section are the following:

- (i) The Legendrian knot K_0 in $\mathbf{R}^3$ whose front projection is a simple closed curve with two cusps has the invariants $tb(K_0) = -1$ and $r(K_0) = 0$.
- (ii) By approximating the front projection of K_0 by a front with an additional zigzag (a left and a right cusp, oriented up or down) and no self-crossings, we can decrease the Thurston-Bennequin invariant by 1 (i.e. add a negative twist to the contact framing) and decrease or increase the rotation number by 1. Hence, by adding n zigzags we can realise $tb = -n - 1$ and any of the $n + 1$ different rotation numbers

$$-n, -n + 2, -n + 4, \dots, n - 2, n,$$

i.e. all values in steps by 2 in the range allowed by the Thurston-Bennequin inequality.

Another useful fact that can easily be proved using the concept of front projections is that any knot $p(t) = (x(t), y(t), z(t))$ in $\mathbf{R}^3$ can be C^0 -approximated by a Legendrian knot: simply approximate the plane curve $(y(t), z(t))$ by a front (with many zigzags) whose slope approximates $-x(t)$. The same is true for knots in arbitrary contact manifolds.

5 Equivalent contact surgeries

The strategy for proving Theorem 1 is as follows. Let (M, ξ) be given. By the topological surgery presentation theorem of Lickorish [18] and Wallace [24], the manifold M can be obtained by performing integer surgery along a link in S^3 . Thus, conversely, we can recover S^3 by integer surgery along a link in M . By the remark at the end of the preceding section, we may assume this to be a Legendrian link, so that we can perform the surgeries as contact integer surgeries. The final product of these surgeries will be S^3 with some contact structure ξ'_0 .

In the present section we show that all these contact integer surgeries may be replaced by contact (± 1) -surgeries along a different link. In the next section we show how to transform (S^3, ξ'_0) into (S^3, ξ_0) by further contact (± 1) -surgeries. Notice that the surgery curves may be moved about by contact isotopies (which exist in abundance due to their constructibility via contact Hamiltonians); this allows to replace an iteration of contact surgeries by a simultaneous contact surgery (without altering the surgery coefficients) along a suitable link.

Proposition 3. *Any contact r -surgery with $r < 0$ can be obtained by a sequence of contact (-1) -surgeries.*

Proof. We want to show that a contact r -surgery, $r < 0$, along a Legendrian knot K may be replaced by a sequence of contact (-1) -surgeries along suitable Legendrian knots inside a tubular neighbourhood νK of K . This would seem to suggest that one should identify νK with $S^1 \times D^2$ using the contact framing of K . For our inductive procedure, however, it turns out to be more opportune to twist this framing by -1 .

Thus, we set $N_0 = S^1 \times D^2$ and write μ_0 for its meridian and λ_0 for the longitude determined by setting φ equal to a constant (in the usual coordinates (θ, r, φ) on $S^1 \times D^2$). Recall the following result [17, Thm. 8.2], cf. [16, Prop. 4.3] (which is the reason why contact $(1/k)$ -surgery is uniquely defined, see [2, Prop. 7]):

Proposition 4. *For any integer k (including 0) there exists a unique (up to isotopy fixed at the boundary) tight contact structure on $S^1 \times D^2$ with a fixed convex boundary with $\#\Gamma_{\partial(S^1 \times D^2)} = 2$ and slope $s(\partial(S^1 \times D^2)) = 1/k$.*

Let ξ be the essentially unique tight contact structure on N_0 with $\#\Gamma_{\partial N_0} = 2$ and boundary slope -1 . In other words, the longitude determined by the contact framing is $\lambda_0 - \mu_0$. Thus, if μ denotes the meridian of νK and λ the longitude of νK determined by the contact framing, then the identification of N_0 with νK is given by

$$\mu_0 \longmapsto \mu, \quad \lambda_0 \longmapsto \mu + \lambda,$$

so that $\lambda_0 - \mu_0 \mapsto \lambda$.

If we let the meridian correspond to the first coordinate direction in $\mathbf{R}^2/\mathbf{Z}^2$, and the longitude to the second, then the above identification is described by the matrix $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Now let K_1 be a Legendrian knot in N_0 , isotopic to the spine $S^1 \times \{0\}$, and with boundary slope $1/(r_1+1)$, $r_1+1 \leq -1$. That is, K_1 has a tubular neighbourhood $N_1 \subset N_0$ (with convex boundary ∂N_1 with $\#\Gamma_{\partial N_1} = 2$) such that, under the diffeomorphism of ∂N_1 with ∂N_0 determined by the isotopy of K_1 with $S^1 \times \{0\}$, the longitude of ∂N_1 determined by the contact framing is equal to $\lambda_0 + (r_1 + 1)\mu_0$ (and meridian still equal to μ_0). For $r_1 + 1 \leq -1$ this is exactly what remark (ii) in the preceding section allows us to do.

The subscript $i = 0, 1$ in N_i here is of course just a counter and does not refer to the radius of these solid tori. The same applies to all subsequent occurrences of the notation ' N_i '. There should be little grounds for confusion with the notation in Section 2 in the definition of contact r -surgery.

Now perform a contact (-1) -surgery on K_1 . That is, we cut out N_1 and glue back a solid torus N'_1 by sending its meridian μ_1 to

$$\mu_0 - (\lambda_0 + (r_1 + 1)\mu_0) = -r_1\mu_0 - \lambda_0.$$

We may choose a longitude λ_1 for N'_1 such that the gluing is described by the matrix $\begin{pmatrix} -r_1 & 1 \\ -1 & 0 \end{pmatrix} \in \text{SL}(2, \mathbf{Z})$ with respect to (μ_1, λ_1) and (μ_0, λ_0) (that is, λ_1 maps to μ_0). Notice that the surgered N_0 is still a solid torus: after cutting out N_1 we have a torus shell $T^2 \times [0, 1]$, and we glue a solid torus N'_1 to one of its boundary components.

Observe that the curve $\lambda_1 - \mu_1$ on $\partial N'_1$ is mapped to a dividing curve

$$\lambda_0 + (r_1 + 1)\mu_0$$

on ∂N_1 . This determines the extension of the contact structure over N'_1 ; it is the unique tight contact structure for which $\partial N'_1$ is convex with $\#\Gamma_{\partial N'_1} = 2$ and $s(\partial N'_1) = -1$.

We now let N'_1 take the role of N_0 . That is, we choose a Legendrian knot isotopic to the spine of N'_1 and with boundary slope $1/(r_2 + 1)$ (expressed in terms of (μ_1, λ_1)), where $r_2 + 1 \leq -1$, and we perform contact (-1) -surgery on this knot. Continuing this process, we see that by a sequence of (-1) -surgeries (say n of them) we can realise a topological r -surgery, where $r = p/q$ is related to the integers $r_1, \dots, r_n \leq -2$ by

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} -r_1 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} -r_2 & 1 \\ -1 & 0 \end{pmatrix} \cdots \begin{pmatrix} -r_n & 1 \\ -1 & 0 \end{pmatrix} = \begin{pmatrix} p & p' \\ q & q' \end{pmatrix}.$$

From this one finds that p/q has the continued fraction expansion

$$r_1 + 1 - \frac{1}{r_2 - \frac{1}{\dots - \frac{1}{r_n}}},$$

see [22, Thm. 7.1.1], for instance. We abbreviate this continued fraction as $[r_1 + 1, r_2, \dots, r_n]$. The following lemma we leave as a simple exercise (use the Euclidean algorithm with negative remainders).

Lemma 5. *Any rational number $r < 0$ can be written in the form*

$$r = [r_1 + 1, r_2, \dots, r_n]$$

with $r_1, \dots, r_n \leq -2$.

The next thing to notice is that the topological r -surgery we obtain in this way is actually a contact r -surgery. For this we only need to show that we still have a tight contact structure on the surgered N_0 (in general, although contact r -surgery is defined by choosing a tight extension over the solid torus to be glued in, the resulting contact structure on the surgered manifold need not be tight). But this follows from the fact that contact (-1) -surgeries correspond to symplectic handlebody surgeries. Furthermore, N_0 has a contact embedding into (S^3, ξ_0) , which is symplectically filled by the standard 4-ball B^4 . So we can realise the surgered N_0 as a subset of the manifold obtained as the new boundary after attaching n symplectic 2-handles to B^4 along $S^3 = \partial B^4$. As a symplectically fillable contact manifold, it is tight. See [6] for more details on the notions used in this paragraph.

It remains to show that *any* contact r -surgery can be performed in this way. The net result of performing the sequence of (-1) -surgeries inside N_0 is a solid torus with a tight contact structure and with convex boundary ∂N_0 whose dividing set has two components (for the boundary has remained unchanged). The slope of the dividing set has changed, however. To determine this slope, we need to find the curve on $\partial N'_n$ (the boundary of the last solid torus to be glued in) that maps to $\lambda_0 - \mu_0$ under the successive gluings. That is, $s(\partial N_0) = y/x$, where x and y are determined by

$$\begin{pmatrix} -r_1 & 1 \\ -1 & 0 \end{pmatrix} \cdots \begin{pmatrix} -r_n & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}.$$

In analogy with the considerations above one finds $y/x = [r_n, \dots, r_2, r_1 + 1]$. Notice that

$$[r_n, \dots, r_2, r_1 + 1] = [r_n, \dots, r_{k+1}, r_k + 1],$$

where k is the smallest index for which $r_k < -2$ (if all r_i are equal to -2 , then the continued fraction equals -1). As proved by Honda [16, Thm. 2.3], cf. [12], the number of tight contact structures on the solid torus with this convex boundary is equal to

$$|(r_n + 1) \cdots (r_1 + 1)|$$

(beware that Honda's notation differs slightly from ours). We claim that this is exactly the number of tight contact structures we can produce by choosing our surgery curves $K_1, \dots, K_n$ suitably. Indeed, we had fixed $tb(K_i) = r_i + 1 \leq -1$ (where we regard K_i as a Legendrian knot in $N'_{i-1} \equiv N_0 \subset (S^3, \xi_0)$), and then by remark (ii) in the preceding section we have a choice of $|r_i + 1|$ different rotation numbers. One now argues as in the proof of [16, Prop. 4.18] (building on a result of Lisca and Matić) that the resulting contact structures are pairwise distinct.

This completes the proof of Proposition 3.

As a direct consequence of this proof we see that our problem of replacing integer surgeries by (± 1) -surgeries has a simple solution for negative surgery coefficients.

Corollary 6. *Any contact n -surgery with $n \in \mathbf{Z}^-$ can be replaced by a suitable (-1) -surgery.*

Next we deal with positive integer surgery coefficients. Again we begin by proving a more general statement about surgery with positive rational coefficient.

Proposition 7. *Any contact r -surgery with $r > 0$ can be obtained by a contact $(1/k)$ -surgery with some positive integer k , followed by a suitable contact r' -surgery with $r' < 0$.*

Proof. Write $r = p/q$ with p, q coprime positive integers. Choose integers p', q' such that $pq' - qp' = 1$. Then, topologically, r -surgery along a Legendrian knot K is defined by cutting out a tubular neighbourhood νK and gluing back a solid torus N_0 with the attaching map

$$\mu_0 \longmapsto p\mu + q\lambda, \quad \lambda_0 \longmapsto p'\mu + q'\lambda,$$

where the notation is as in the proof of Proposition 3; in particular, λ denotes the longitude determined by the contact framing of K .

This means that $-p'\mu_0 + p\lambda_0$ is glued to the dividing curve λ . So the possible contact r -surgeries are determined by the tight contact structures on N_0 with convex boundary satisfying $\#\Gamma_{\partial N_0} = 2$ and $s(\partial N_0) = -p/p'$.

Now choose a positive integer k such that $q - kp < 0$. We can perform a $(1/k)$ -surgery on K by cutting out νK and gluing in N_0 , with gluing map

$$\mu_0 \longmapsto \mu + k\lambda, \quad \lambda_0 \longmapsto \lambda.$$

The unique contact $(1/k)$ -surgery is defined by the tight contact structure on N_0 with convex boundary satisfying $\#\Gamma_{\partial N_0} = 2$ and $s(\partial N_0) = \infty$ (in terms of (μ_0, λ_0)), in other words, by taking N_0 to be the standard neighbourhood of a Legendrian knot.

Next perform r' -surgery along the spine of N_0 , with $r' = p/(q - kp)$. That is, cut out a tubular neighbourhood N_1 of the spine of N_0 and glue back a solid torus N'_1 with the attaching map

$$\mu_1 \longmapsto p\mu_0 + (q - kp)\lambda_0, \quad \lambda_1 \longmapsto p'\mu_0 + (q' - kp')\lambda_0.$$

Since

$$\begin{pmatrix} 1 & 0 \\ k & 1 \end{pmatrix} \begin{pmatrix} p & p' \\ q - kp & q' - kp' \end{pmatrix} = \begin{pmatrix} p & p' \\ q & q' \end{pmatrix},$$

the net result of these two surgeries is the desired r -surgery.

We see that $-p'\mu_1 + p\lambda_1$ is glued to λ_0 , so again the possible contact surgeries are the ones corresponding to tight contact structures on the solid torus with boundary slope $-p/p'$. Moreover, since rational contact surgery with negative coefficient can be effected by a sequence of contact (-1) -surgeries, i.e. symplectic handlebody surgeries, the result of performing a contact r' -surgery on N_0 (which can be realised on $N_0 \subset (S^3, \xi_0)$, for instance) will be a tight contact structure on all of N_0 . So the different choices correspond exactly to the different contact r -surgeries along K .

Again we can specialise this to contact surgeries with positive integer coefficients.

Corollary 8. *Any contact n -surgery with $n \in \mathbf{Z}^+$, $n \geq 2$, is equivalent to a contact $(+1)$ -surgery followed by a contact $(n/(1-n))$ -surgery, and hence (by Proposition 3) to a contact $(+1)$ -surgery followed by a sequence of contact (-1) -surgeries.*

The only case left open is that of a contact 0-surgery. We claim that the result of any contact 0-surgery is the same as that of a suitable contact $(+1)$ -surgery. We defer the proof of this fact to the next section, because it can best be dealt with in the context of Lutz twists. Taking this statement about 0-surgery for granted, we have proved that by performing contact (± 1) -surgeries along a Legendrian link in M , we can obtain S^3 with some contact structure.

6 Lutz twists and contact surgeries

In order to prove Theorem 1 it remains to show that, given any contact structure ξ'_0 on S^3 , one can find a Legendrian link in S^3 such that contact (± 1) -surgery along this link produces (S^3, ξ_0) . For this it is sufficient to observe that (S^3, ξ'_0) can be obtained from (S^3, ξ_0) by suitable so-called *Lutz twists*, and that any Lutz twist is equivalent to certain integer surgeries. We shall now elaborate on these points.

First recall the concept of a Lutz twist. Let K be any knot in a contact 3-manifold (M, ξ) that is transverse to ξ . Then K has a tubular neighbourhood $S^1 \times D_{r_0}^2 \subset M$, where $D_{r_0}^2$ denotes the open 2-disc of radius r_0 and $K \equiv S^1 \times \{0\}$, on which ξ is given by $d\theta + r^2 d\varphi$ in the coordinates (θ, r, φ) used before, cf. [21]. By a diffeomorphism of the solid torus $S^1 \times D_{r_0}^2$ given by Dehn twists along the meridian (which amounts to a different choice of longitude) and rescaling the r -coordinate we may assume that $r_0 > 1$.

A *simple Lutz twist* along K is defined by replacing ξ on $S^1 \times D_{r_0}^2$ by the contact structure $\beta_0 = 0$, where $\beta_0 = h_1(r) d\theta + h_2(r) d\varphi$ is defined as in Section 2, subject to the following conditions (for some small $\varepsilon > 0$):

- $h_1(r) \equiv -1$ and $h_2(r) \equiv -r^2$ for $r < \varepsilon$,

- $h_1(r) \equiv 1$ and $h_2(r) = r^2$ for $r > 1 - \varepsilon$,
- $(h_1(r), h_2(r))$ does *not* complete a full turn around $(0, 0)$ as r goes from 0 to r_0 .

Such a simple Lutz twist does not change the topology of the underlying manifold, but it does, in general, change the homotopy type of the contact structure (as a 2-plane field).

A *full Lutz twist* is defined by the conditions

- $h_1(r) \equiv 1$ and $h_2(r) = r^2$ for $r < \varepsilon$ and $r > 1 - \varepsilon$,
- $(h_1(r), h_2(r))$ completes exactly one full twist around $(0, 0)$.

A full Lutz twist does not change the homotopy type of the contact structure as a 2-plane field, nor the topology of the underlying manifold.

Notice that the disc $\{\theta_0\} \times D_{r'_0}^2$, where $r'_0 > 0$ is chosen such that $h_2(r'_0) = 0$, is an overtwisted disc (after a small perturbation). In the case of a simple Lutz twist, r'_0 is unique; in the case of a full Lutz twist there are two choices.

Homotopy classes of (cooriented) 2-plane fields on S^3 are classified by the homotopy group $\pi_3(S^2) \cong \mathbf{Z}$. As shown by Lutz [19], cf. [20] (again, the full details are folklore), each of these homotopy classes contains a contact structure, obtained by suitable Lutz twists from (S^3, ξ_0) . Moreover, Eliashberg [5] has given a complete classification (up to isotopy) of contact structures on S^3 : Each homotopy class contains exactly one overtwisted contact structure, and the only tight contact structure is the standard one ξ_0 . The overtwisted contact structure that is homotopic to ξ_0 as a 2-plane field can be obtained from ξ_0 by a full Lutz twist. In conclusion, every contact structure on S^3 can be obtained from ξ_0 by Lutz twists.

Proposition 9. *A simple Lutz twist is equivalent to two contact (+1)–surgeries.*

Proof. (cf. [7, Prop. 4.3]) Consider $N_0 = S^1 \times D_{r_0}^2$ with $r_0 > 1$ and contact structure ζ_0 given by $h_1(r) d\theta + h_2(r) d\varphi = 0$, with h_1, h_2 as described above for a simple Lutz twist.

Let r_1 be the unique radius for which $h_1(r_1) = -h_2(r_1) > 0$. Then along the 2-torus $T_{r_1} = \{r = r_1\}$ the contact structure ζ_0 is given by $d\theta - d\varphi = 0$, so the characteristic foliation on T_{r_1} is linear of slope $d\theta/d\varphi = 1$. As r goes from 0 to r_1 , the slope of the characteristic foliation on T_r decreases monotonically from 0 to $-\infty$, and then further from ∞ to 1. This shows that the restriction of ζ_0 to $N_1 = S^1 \times D_{r_1}^2$ is tight, for by Dehn twists along the meridian and rescaling the r -coordinate we can embed this into the solid torus with its standard tight contact structure $d\theta + r^2 d\varphi = 0$.

Now perturb T_{r_1} into a convex torus of slope 1 as described at the end of Section 2. Then the restriction of ζ_0 to the perturbed N_1 is the tight contact structure uniquely determined by this convex boundary and may be thought of as a standard neighbourhood of its Legendrian spine K_1 . Notice that the contact framing of K_1 is given by $\lambda_0 + \mu_0$.

As a next step, we perform contact (−1)–surgery along K_1 . That is, we cut out N_1 and glue back a solid torus N'_1 with the attaching map

$$\mu_1 \longmapsto \mu_0 - (\lambda_0 + \mu_0) = -\lambda_0, \quad \lambda_1 \longmapsto \mu_0.$$

This means that $\lambda_1 - \mu_1$ is mapped to the dividing curve $\lambda_0 + \mu_0$, so N'_1 carries the unique tight contact structure with convex boundary of slope -1 .

We now claim that the net result of performing a simple Lutz twist and the described contact (-1) -surgery is a contact $(+1)$ -surgery. To prove this, perturb the 2-torus $T_1 = \{r = 1\}$ into a convex torus with dividing set of slope -1 with respect to (μ_0, λ_0) (and two components, as usual). Consider the thickened torus $T^2 \times [r_1, 1]$ with boundaries the perturbed T_{r_1} and T_1 . The restriction of ζ_0 to this thickened torus is tight (after a diffeomorphism of the thickened torus it is seen to embed into the standard tight structure on $S^1 \times D^2$), the boundaries are convex, and ζ_0 is *minimally twisting* in the sense of [16, Section 2.2.1], that is, the contact structure twists minimally in radial direction to satisfy the boundary conditions. Notice that with respect to (μ_1, λ_1) the slope of T_1 is $+1$, since $\lambda_1 + \mu_1$ is glued to $\mu_0 - \lambda_0$. Moreover, with respect to (μ_1, λ_1) the slope twists from -1 on T_{r_1} to $+1$ on T_1 by passing via $\pm\infty$ rather than 0 (λ_1 is glued to μ_0).

Let r'_1 be the radius determined by $h_1(r'_1) = h_2(r'_1) < 0$. We then see that $T^2 \times [r_1, 1]$ is contactomorphic to $T^2 \times [r'_1, r_1]$ (either equipped with the restriction of ζ_0 , and with perturbed boundaries). Furthermore, by Proposition 4 we know that N'_1 is contactomorphic to the perturbed $S^1 \times D^2_{r'_1}$. We conclude that $N'_1 \cup T^2 \times [r_1, 1]$ is contactomorphic to the perturbed N_1 , i.e. the solid torus with the essentially unique tight contact structure with convex boundary of slope $+1$.

The perturbed T_1 has slope -1 with respect to (μ_0, λ_0) , so it may be regarded as the boundary of a standard tubular neighbourhood of a Legendrian knot with contact framing $\lambda_0 - \mu_0$. So the result of performing a Lutz twist and a (-1) -surgery as described is to cut out this tubular neighbourhood, glue in a solid torus N'_2 with attaching map that sends μ_2 to $-\lambda_0 = -\mu_0 - (\lambda_0 - \mu_0)$, and extend with the unique tight contact structure on N'_2 determined by this gluing. This amounts to a contact $(+1)$ -surgery.

We conclude that this last contact $(+1)$ -surgery, followed by the contact $(+1)$ -surgery reversing the (-1) -surgery along K_1 , amounts to a simple Lutz twist.

As a more or less immediate corollary of this proof we have the following results. They complete the proof of Theorem 1.

Proposition 10. (i) *A full Lutz twist is equivalent to four contact $(+1)$ -surgeries.*

(ii) *A contact 0-surgery is equivalent to a contact $(+1)$ -surgery.*

Proof. Argue as in the proof of the preceding proposition. For (i), perform a contact (-1) -surgery determined by the torus T_{r_1} , where $h_1(r_1) = -h_2(r_1) < 0$, then a contact (-1) surgery determined by T_{r_2} , where $h_1(r_2) = h_2(r_2) < 0$. This reduces a full Lutz twist to a simple Lutz twist.

For (ii), perform a contact (-1) -surgery as in the case of a simple Lutz twist. This reduces the contact 0-surgery to a trivial surgery. (The relation between the two Legendrian knots corresponding to the contact 0-surgery and $(+1)$ -surgery, respectively, is analogous to that in the case of contact surgeries with different negative integer framings, as discussed in the proof of Proposition 3.)

Acknowledgements.

This research was carried out while F. D. was staying at the Mathematical Institute of Leiden University (then the home of H. G.), supported by a post-doctoral fellowship from NWO (Netherlands Organisation for Scientific Research). The research of F. D. is partially supported by the NSFC (National Science Foundation of China).

References

- [1] V.I. Arnol'd, Lagrange and Legendre cobordisms I, *Funct. Anal. Appl.* **14** (1980), 167–177.
- [2] F. Ding and H. Geiges, Symplectic fillability of tight contact structures on torus bundles, *Algebr. Geom. Topol.* **1** (2001), 153–172.
- [3] Y. Eliashberg, Classification of overtwisted contact structures on 3-manifolds, *Invent. Math.* **98** (1989), 623–637.
- [4] Y. Eliashberg, Topological characterization of Stein manifolds of dimension > 2 , *Internat. J. of Math.* **1** (1990), 29–46.
- [5] Y. Eliashberg, Contact 3-manifolds twenty years since J. Martinet's work, *Ann. Inst. Fourier (Grenoble)* **42** (1992), 165–192.
- [6] J.B. Etnyre, Symplectic convexity in low-dimensional topology, *Topology Appl.* **88** (1998), 3–25.
- [7] J.B. Etnyre and K. Honda, On symplectic cobordisms, *Math. Ann.* **323** (2002), 31–39.
- [8] H. Geiges, A brief history of contact geometry and topology, *Expo. Math.* **19** (2001), 25–53.
- [9] H. Geiges, Contact geometry, in: *Handbook of Differential Geometry*, vol. 2 (F.J.E. Dillen, L.C.A. Verstraelen, eds.), to appear.
(available from <http://www.mi.uni-koeln.de/~geiges/publications.html>)
- [10] E. Giroux, Convexit  en topologie de contact, *Comment. Math. Helv.* **66** (1991), 637–677.
- [11] E. Giroux, Une infinit  de structures de contact tendues sur une infinit  de vari t s, *Invent. Math.* **135** (1999), 789–802.
- [12] E. Giroux, Structures de contact en dimension trois et bifurcations des feuilletages de surfaces, *Invent. Math.* **141** (2000), 615–689.

- [13] E. Giroux, Géométrie de contact: de la dimension trois vers les dimensions supérieures, in: *Proceedings of the International Congress of Mathematicians* (Beijing, 2002) vol. II, Higher Education Press, Beijing (2002), 405–414.
- [14] E. Giroux, Structures de contact, livres ouverts et tresses fermées, in preparation.
- [15] R.E. Gompf, Handlebody construction of Stein surfaces, *Annals of Math. (2)* **148** (1998), 619–693.
- [16] K. Honda, On the classification of tight contact structures I, *Geom. Topol.* **4** (2000), 309–368.
- [17] Y. Kanda, The classification of tight contact structures on the 3–torus, *Comm. Anal. Geom.* **5** (1997), 413–438.
- [18] W.B.R. Lickorish, A representation of orientable combinatorial 3–manifolds, *Annals of Math. (2)* **76** (1962), 531–540.
- [19] R. Lutz, Sur l’existence de certaines formes différentielles remarquables sur la sphère S^3 , *C. R. Acad. Sci. Paris Sér. A-B* **270** (1970), A1597–A1599.
- [20] R. Lutz, Structures de contact sur les fibrés principaux en cercles de dimension trois, *Ann. Inst. Fourier (Grenoble)* **27**, no. 3 (1977), 1–15.
- [21] J. Martinet, Formes de contact sur les variétés de dimension 3, in: *Proceedings of the Liverpool Singularities Symposium II*, Lecture Notes in Math. **209**, Springer-Verlag, Berlin, 1971, pp. 142–163.
- [22] H.E. Rose, *A Course in Number Theory*, Oxford University Press, 1988.
- [23] A. Stipsicz, private communication.
- [24] A.H. Wallace, Modifications and cobounding manifolds, *Canad. J. Math.* **12** (1960), 503–528.
- [25] A. Weinstein, Contact surgery and symplectic handlebodies, *Hokkaido Math. J.* **20** (1991), 241–251.

Mathematisches Institut, Universität zu Köln, Weyertal 86–90, 50931 Köln, Germany;
 e-mail: geiges@math.uni-koeln.de
 Department of Mathematics, Peking University, Beijing 100871, P. R. China, e-mail:
 dingfan@math.pku.edu.cn